

Contribution ID: 48 Type: not specified

Development of a new B-Physics trigger for the ATLAS Detector at CERN (student talk)

Saturday, 17 February 2018 11:45 (15 minutes)

A novel B-Physics trigger for the ATLAS detector is being developed to enable lepton universality studies by selecting $B^0 \to K^*e^+e^-$ events and complementing the existing $B^0 \to K^*\mu^+\mu^-$ trigger. In the Standard Model, lepton universality refers to the fact that the electroweak couplings of the leptons to the gauge bosons is independent of the lepton flavour. A particular sensitive probe for studying lepton universality is by measuring the ratio $(R_{K^{*0}})$ of the branching fractions of $B^0 \to K^{*0}\mu^+\mu^-$ and $B^0 \to K^{*0}e^+e^-$ decay processes. The value of $R_{K^{*0}}$ is expected to be close to unity in the Standard Model. However, recent analyses have shown an intriguing deviation from the expected value of $R_{K^{*0}}$. The development of the $B^0 \to K^{*e}e^+e^-$ trigger is critical to the measurement of the ratio. Summary of ongoing work will be presented in the talk.

Primary authors: Dr RUSSELL, Heather (McGill University); Ms SAHA, Shreya (McGill University); Prof.

ROBERTSON, Steven (McGill University)

Presenter: Ms SAHA, Shreya (McGill University)

Session Classification: Session #4