

Production of ^{230}Pa from proton-irradiated thorium and developing $^{230}\text{Pa}/^{230}\text{U}/^{226}\text{Th}$ tandem generator

Targeted alpha-therapy of various oncology diseases is based on the coupling of alpha-emitting radionuclides to tumour-selective carrier molecules. ^{230}U ($T_{1/2}=20.8$ d) is accumulated via decay of ^{230}Pa and has strong potential for alpha-therapy due to 5 emitting alpha - particles with total energy 33.5 MeV. It can be used directly or as a parent of ^{226}Th in a generator system.

A prospective way for production of $^{230}\text{Pa}/^{230}\text{U}$ is irradiation of natural thorium with medium-energy protons. Curie amounts of ^{230}Pa can be generated in one irradiation run together with other useful alpha-emitters ^{225}Ac and ^{223}Ra .

After dissolving Th-target in 6M nitric acid with the addition of catalytic amounts of hydrofluoric acid Pa was isolated by liquid extraction with octanol. The water phase may be further used for isolation of ^{225}Ac and ^{223}Ra according the method, proposed in [1, 2]. Namely, the most part of thorium was removed by liquid extraction with di(2-ethylhexyl)phosphoric acid in toluene. Organic phase was kept for accumulation ^{223}Ra from parent ^{227}Th . ^{225}Ac and ^{223}Ra were concentrated from aqueous phase and then separated and purified by extraction chromatography. After re-extraction ^{230}Pa was purified on silica gel using different oxalic acid solutions. Up to 85% of ^{230}Pa with radionuclidic purity >99% was recovered and stored for ^{230}U accumulation.

Basing on distribution coefficients ^{230}U was separated from ^{230}Pa in diluted nitric acid solution on DGA resin. Obtained ^{230}U may be used for development of $^{230}\text{U}/^{226}\text{Th}$ -generator. Distribution coefficients for U and Th were obtained for a wide range of extraction chromatographic resins. TEVA (mixture of trioctyl and tridecyl methyl ammonium chloride as an extractant) and WBEC (based on a mixture of tertiary octyl and decylamines) resins are most suitable for this purpose.

REFERENCES

1. R.A. Aliev, S.V. Ermolaev, A.N. Vasiliev, V.S. Ostapenko, E.V. Lapshina, B.L. Zhuikov, N.V. Zakharov, V.V. Pozdeev, V.M. Kokhanyuk, B.F. Myasoedov, S.N. Kalmykov. Isolation of medicine-applicable actinium-225 from thorium targets irradiated by medium-energy protons. Solvent Extraction and Ion Exchange, 2014, v. 32, p. 468-477.
2. A.N. Vasiliev, V.S. Ostapenko, E.V. Lapshina, S.V. Ermolaev, S.S. Danilov, B.L. Zhuikov and S. N. Kalmykov. Recovery of Ra-223 from natural thorium irradiated by protons. Radiochimica Acta, 2016, v. 104, N.8, p. 539-547.

Email Address

lapshina@inr.ru

Presentation Type

Contributed Oral

Primary author: Dr LAPSHINA, Elena (Institute for Nuclear Research of Russian Academy of Sciences)

Co-authors: Dr VASILIEV, Aleksandr (Institute for Nuclear Research of Russian Academy of Sciences); Dr ZHUIKOV, Boris (Institute for Nuclear Research of Russian Academy of Sciences); Dr ERMOLAEV, Stanislav (Institute for Nuclear Research of Russian Academy of Sciences); Ms OSTAPENKO, Valentina (Institute for Nuclear Research of Russian Academy of Sciences)

Presenter: Dr LAPSHINA, Elena (Institute for Nuclear Research of Russian Academy of Sciences)