



Contribution ID: 7

Type: Nuclear and Particle Physics

Mirror Symmetry in the $f_{1/2}$ Shell Below ^{56}Ni , Excited States and Electromagnetic Transition Rates in ^{55}Ni and ^{55}Co

Tuesday, 15 February 2022 08:48 (12 minutes)

Experiment S1758 aims to explore the charge dependence of the strong nuclear interaction by probing ^{55}Ni and ^{55}Co near the *doubly magic* ^{56}Ni . This will be achieved by impinging beams of radioactive ^{20}Na and stable ^{20}Ne upon ^{40}Ca targets to produce ^{55}Ni and ^{55}Co , respectively. Charged particles and γ -rays will be detected by combining TRIUMF-ISAC Gamma-Ray Escape Suppressed Spectrometer (TIGRESS), the TIGRESS Integrated Plunger (TIP) and the CsI Ball. This trio allows for a higher degree of sensitivity when in unison. Data analysis will involve: transition rate reconstruction using the Doppler-Shift Attenuation Method (DSAM), Doppler-shift lineshape profile extraction from Monte Carlo simulations via the GEANT4 framework, and lifetime extraction from minimizing a χ^2 goodness-of-fit between the measured and simulated lineshapes. The results will paint a clearer picture of the charge dependence of the strong nuclear interaction.

email address

tschilba@sfu.ca

Please select: Experiment or Theory

Experiment

Primary author: SCHILBACH, Tyson (Simon Fraser University Physics Department)

Presenter: SCHILBACH, Tyson (Simon Fraser University Physics Department)

Session Classification: Nuclear Physics