

## Update on the *hep* solar neutrino limit from the three-phase SNO dataset

Thursday, 1 November 2018 18:00 (3 minutes)

The spectrum of solar neutrinos from the *pp* chain has been studied in depth by a variety of underground detectors. However, neutrinos from the *hep* reaction ( ${}^3\text{He} + p^+ \rightarrow {}^4\text{He} + e^+ + \nu_e$ ) remain unobserved due to the small theoretical branching ratio ( $2 \times 10^{-7}$  per *pp* termination). The SNO detector has a unique sensitivity to neutrino energies above the  ${}^8\text{B}$  spectrum endpoint ( $\sim 15$  MeV) through the *hep* spectrum endpoint ( $\sim 18.8$  MeV) due to the charged current interaction on deuterium, which allows a more precise extraction of the underlying neutrino energy spectrum. The SNO collaboration previously published a world-leading limit in 2006 only using the first heavy water phase with 306.4 days of data. An updated status report on the analysis of the *hep* neutrino spectrum from all three phases of SNO (1170.2 days) will be presented in this poster.

**Primary author:** Dr GILJE, Karin (University of Alberta)

**Co-author:** Dr MASTBAUM, Andy (University of Chicago)

**Presenter:** Dr GILJE, Karin (University of Alberta)

**Session Classification:** Poster session and Reception