



Contribution ID: 106

Type: **Presentation with support**

Interaction Cross Sections and Neutron Skin Thickness of Ni Isotopes

Monday, 19 August 2024 15:35 (15 minutes)

Since nuclear matter is composed of two Fermi particles, protons and neutrons, the equation of state of nuclear matter has a term that depends on the density difference between the two, which is called the symmetry energy. From previous studies, it is known that the first-order density dependence of the symmetry energy is closely related to the thickness of the neutron skin [1].

In this study, interaction cross sections σ_I and charge changing cross sections σ_{CC} for $^{58-77}\text{Ni}$ on a carbon target at 260 MeV/nucleon have been measured to derive matter radii and charge radii respectively. Recently, the charge radii of Ni isotopes up to mass number 70 were measured by isotope shift method [2]. In order to derive the neutron skin thickness in the more neutron-rich region, we attempted to derive the charge radii from charge changing cross section measurements. The experiment was performed at the Radioactive Isotope Beam Factory (RIBF) at RIKEN by using the BigRIPS fragment separator.

In this presentation, we'll report the matter radii and charge radii derived from the experimental cross sections using Glauber calculations. Also, in the region where the charge radii are known, from $A = 58$ to 70, we'll discuss the neutron skin thickness of Ni isotopes, which is obtained by the present data combined with known charge radii. On the other hand, in the neutron-rich region with $A \geq 71$, we'll discuss the neutron skin thickness by combining present σ_I and σ_{CC} data.

References

- [1] M. Centelles et al., Phys. Rev. Lett. 102 (2009) 122502.
- [2] S. Malbrunot-Ettenauer et al., Phys. Rev. Lett. 128 (2022) 022502

Funding Agency

Email Address

fukutome@ne.phys.sci.osaka-u.ac.jp

Presenter if not the submitter of this abstract

Primary authors: Prof. PROCHAZKA, A. (GSI); Mr HOMMA, Akira (Niigata Univ.); Prof. OZAWA, Akira (Univ. of Tsukuba); Mr YANO, Asahi (Univ. of Tsukuba); Prof. SCHEIDENBERGER, C. (GSI); Mr FUKUSHIMA, Chihaya (Tokyo City Univ.); Prof. NISHIMURA, Daiki (Tokyo City Univ.); Prof. NAGAE, Daisuke (RIKEN); Prof. MIYATA, Eri (Niigata Univ.); Mr TAKAYAMA, Gen (Osaka Univ.); Prof. GEISSEL, H. (GSI); Prof. SAKURAI, Hiroyoshi (RIKEN, Univ. of Tokyo); Prof. TAKEDA, Hiroyuki (RIKEN); Prof. TANIHATA, Isao (Osaka Univ. RCNP); Prof. BEHR, K.-H. (GSI); Prof. MATSUTA, Kensaku (Osaka Univ.); Prof. KUSAKA, Kensuke (RIKEN); Prof. ITABASHI, Kenta (RIKEN); Prof. YOSHIDA, Koichi (RIKEN); Prof. OHTAKE, Masao (RIKEN); Prof. TANAKA,

Masaomi (Kyusyu Univ.); Prof. TAKECHI, Maya (Niigata Univ.); Ms AMITANI, Mei (Tokyo City Univ.); FUKUTOME, Miki; Prof. FUKUDA, Mistunori (Osaka Univ.); Mr MIHARA, Mototsugu (Osaka Univ.); Prof. FUKUDA, Naoki (RIKEN); Prof. INABE, Naoto (RIKEN); Mr NOGUCHI, Norihide (Niigata Univ.); Prof. H. J, Ong (Osaka Univ. RCNP); Mr TAGUCHI, Ryo (Osaka Univ.); Prof. BAGCHI, S. (GSI); Prof. SUZUKI, Shinji (Univ. of Tsukuba); Mr ISHITANI, Soshi (Osaka Univ.); Prof. OHTSUBO, Takashi (Niigata Univ.); Mr SUGISAKI, Takato (Osaka Univ.); Prof. YAMAGUCHI, Takayuki (Saitama Univ.); Prof. SUZUKI, Takeshi (Saitama Univ.); Prof. IZUMIKAWA, Takuji (Niigata Univ. RI); Prof. MORIGUCHI, Tetsuaki (Univ. of Tsukuba); Prof. SUMIKAMA, Toshiyuki (RIKEN); Prof. SHIMIZU, Yohei (RIKEN); Prof. YANAGISAWA, Yoshiyuki (RIKEN); Ms TAZAWA, Yuki (Niigata Univ.); Mr KOBAYASHI, Yukiya (Niigata Univ.); Prof. KORKULU, Z. (RIKEN)

Presenter: FUKUTOME, Miki

Session Classification: Nuclear Reactions I

Track Classification: Nuclear Structure from Collisions