


# Full Event Interpretation at Belle II

## Characterization and Comparison of ML-Based Semileptonic B Tagging



Pranav Prasad

Supervised by: Prof. Bob Kowalewski & Dr. Philipp Horak

February 13, 2026

Department of Physics and Astronomy

University of Victoria | Belle II Experiment



University  
of Victoria



# Belle II at SuperKEKB

## SuperKEKB Accelerator

- $e^+e^-$  collider in Tsukuba, Japan
- Operates at  $\Upsilon(4S)$  (10.58 GeV)
- Produces 2  $B$  mesons pairs nearly at rest




## Belle II Experiment

- General-purpose flavor physics detector
- Precision studies of  $B$ ,  $D$ , and  $\tau$  decays
- Clean  $B\bar{B}$  factory: full reconstruction

Process	$\sigma$ [nb]
$B\bar{B}$	1.1
$q\bar{q}(u, d, s, c)$	4.7
$\tau^+\tau^-$	0.92
<b>Total Hadronic</b>	<b>5.8</b>


Cross sections



# Missing Energy: Why We Use Tagging

## The Challenge: Invisible Neutrinos

- Many interesting  $B$  decays have neutrinos in the final state. ( $B \rightarrow K\nu\bar{\nu}$ ,  $B \rightarrow D^{(*)}l\bar{\nu}$ , etc.)
- Initial state precisely known
- No other particles in the event apart from the two  $B$  mesons
- Directly detected particles from both  $B$  decays overlap in the detector
- **Solution:** Use  $B_{tag}$  to constrain the  $B_{sig}$ .





## 4-momentum Conservation

$$P_{miss} = P_{e^+e^-} - P_{B_{tag}} - P_{visible}$$

# B-Tagging for Missing Energy

- Hadronic Tagging

- $B_{tag} \rightarrow$  hadrons (fully reconstructed)
- **Pro:** Full constraint of  $B_{sig}$  kinematics






# B-Tagging for Missing Energy

- **Hadronic Tagging**

- $B_{tag} \rightarrow$  hadrons (fully reconstructed)
- **Pro:** Full constraint of  $B_{sig}$  kinematics

- **Semileptonic Tagging**


- $B_{tag} \rightarrow D^{(*)} l \nu$  (missing neutrino, like nuclear beta decay!)



# Full Event Interpretation (FEI)

## Full Event

Reconstructs  $\Upsilon(4S)$  event using a bottom up approach.




*Semileptonic Decay*

## ML-Based (FEI)

Hierarchical FastBDT trained on  $> 10,000$  decay chains; generates SigProb to give likelihood of correct reconstruction

## Statistical Gain


Increase in sample size since no hard selection cuts applied but may require calibration



*BDT Training Levels*


# Challenges with FEI Implementation

- High-level features such as BDTs have to be well understood in simulation
- FEI requires careful calibration
- Excited  $D^* \rightarrow D\pi^0/\gamma$  has low mass difference
- Low energy neutrals are often missed in the detector




# Simple Semileptonic Tag

- **Hard cuts** on kinematics and PID (e.g.  $\cos \theta_{B\gamma}$ )
- Apply a  $D^0$  constraint  $m_{D^0} = 1.86\text{GeV}/c^2$
- Allows for events from  $D^* \rightarrow D\pi^0/\gamma$  but does not explicitly reconstruct the  $\pi^0/\gamma$
- **Why Simple?** No multivariate classifier; relies on cuts and vertex fitting
- Ideally should not require corrections and calibration like FEI



# Angular Variables for Semileptonic Tagging




$\cos \theta_{BY}$

$B \rightarrow Y\nu$  where  $Y = D^{(*)}\ell$

$$\cos \theta_{BY} = \frac{2E_B^* E_Y^* - m_B^2 - m_Y^2}{2|\vec{p}_B^*||\vec{p}_Y^*|}$$


**Signal:**  $-1 \leq \cos \theta_{BY} \leq 1$




- **Inside Cone:** Kinematically allowed.
- **Low Tail:**  $m_Y$  underestimated due to missing low energy  $\pi^0/\gamma$ . (Unphysical  $\angle$ )
- **High Tail:**  $E_Y$  overestimated via incorrect association of particles. (Unphysical  $\angle$ )

# FEI vs. Simple Tagging: Decay Topology

- **Simple Tag:** Unreconstructed  $\pi^0/\gamma$  from  $D^{*0} \rightarrow D^0(\pi^0/\gamma)$  shifts the  $D^{*0}\ell$  component toward negative  $\cos\theta_{BY}$  (blue hist).
- **FEI:** Fully reconstructs the decay chain, centering both  $D^0\ell$  and  $D^{*0}\ell$  distributions at zero.





**Simple Tag:** Shifted  $D^{*0}\ell$  component



**FEI Tag:** Centered signal peaks




# (FEI) Tagging- Using Signal Probability

- **Signal Probability:** Multivariate classifier output to select high-purity  $B_{tag}$  candidates. (NOT A TRUE PROBABILITY\*)
- **Selection Tuning:** Tighter cut  $\rightarrow$  higher purity, lower efficiency (High Tail reduced)




# FEI vs. Simple Tagging — $\cos\theta_{BY}$

- **Data–MC:** Out-of-the-box agreement in  $\cos\theta_{BY}$  comparable
- **Background:**  $\sim 20\%$  fewer continuum events (at matched efficiency)
- **Higher Efficiency:** Small increase in background (mostly  $B^0\bar{B}^0$ ) at  $\times 2$  efficiency




# Future Studies

- **BDT Optimization:** Replace hard cuts with BDTs does maximize efficiency
- **Recovery:** FEI recovers  $> 80\%$  of events found via Simple tagging
- **Missing events:** Study non-reconstructed events to identify FEI limitations
- **Hybrid Development:** Co-develop Simple and FEI tags to further improve SL tagging performance



Event overlap (Loose SigProb cut)



- **Performance:** FEI achieves efficiency gains at comparable purity relative to Simple Tag. ( $\sim 0.9\%$  overall tagging efficiency)
- **Data–MC Agreement:** FEI shows improved Data–MC agreement in  $\cos\theta_{BY}$
- **Observation:** Level of agreement is notable, FEI's multivariate structure was expected to require larger corrections and calibration
- **Applications:** Results applicable to all Belle II analyses using semileptonic  $B$  tagging
- **Future Work:** Further optimization and hybrid approaches could yield better performance

# Thank you for listening!



**University  
of Victoria**



Thanks to my supervisors, Prof. Bob Kowalewski and Dr. Philipp Horak, and the UVic Belle II group for their support.

# Backup


# Missing Energy: Why We Use Tagging

## The Challenge: Invisible Neutrinos

- $\Upsilon(4S) \rightarrow B\bar{B}$  kinematics are precisely known.
- $B_{sig}$  with  $\nu$ 's cannot be directly reconstructed.
- **Solution:** Reconstruct  $B_{tag}$  to fully constrain event.

## Key Measurements

- $B \rightarrow K\nu\bar{\nu}$ : Rare SM process; requires tagging to constrain  $B_{sig}$  kinematics (only 1 track).
- $B \rightarrow \tau\nu$ :  $V_{ub}$  measurement; requires tagging for improved efficiency.





$B \rightarrow K\nu\bar{\nu}$  Measurements

## 4-momentum Conservation

$$P_{miss} = P_{e^+e^-} - P_{B_{tag}} - P_{visible}$$

# ROE Variables: $M_{\text{ROE}}$ and $E_{\text{extra}}$



## $M_{\text{ROE}}$

**Invariant mass** of the Rest of Event.

Used to identify background leakage or missing neutrals.

**Target Signal:**  $\approx 0 \text{ GeV}/c^2$

## $E_{\text{extra}}^{\text{ROE}}$




**Calorimeter energy** not associated with  $B_{\text{sig}}$  or  $B_{\text{tag}}$ .

Ensures all energy is accounted for (no "hidden" particles).



**Target Signal:**  $\approx 0 \text{ GeV}$

# FEI vs. Simple Tag: Rest of Event (ROE)

- **Better Resolution:** FEI reduces events with large missing energy ( $> 500$  MeV) by **25%** relative to Simple tagging.
- **Consistency:** Comparable Data–MC agreement maintained across all tags. (Hard to model MC due to beam background and detector effects)



- Data–MC: Comparable agreement in  $M_{\text{ROE}}$  distribution.
- Peak at 135 MeV: From missing  $\pi^0$  in the events possibly from  $B \rightarrow D\pi\ell\nu$



## FEI Selection


- **ROE & Reco**
  - No charged tracks in ROE
  - Only 6  $D$  Common decays used from the FEI (14 in FEI total)
- **Singal Prob**
  - Set to 0.1 to match efficiency of  $D^0\ell$
- **Best Candidate**
  - $B$  SigProb Used for candidate selection

## $D^0\ell$ Selection


- **ROE & Reco**
  - No charged track in ROE
  - 6  $D$  Decay Modes
- **Kinematic Cuts**
  - $\text{InvM} > 2.5$  and  $-3.0 < \cos BY < 1.5$  at steering level
- **Best Candidate**
  - $B$   $\chi$ Prob vertex fit

# FEI vs. Simple Tagging: Decay Topology

- **Simple Tag:** Unreconstructed  $\pi^0/\gamma$  from  $D^{*0} \rightarrow D^0(\pi^0/\gamma)$  shifts the  $D^{*0}\ell$  component toward negative  $\cos\theta_{BY}$  (blue hist).
- **FEI:** Fully reconstructs the decay chain, centering both  $D^0\ell$  and  $D^{*0}\ell$  distributions at zero.





**Simple Tag:** Shifted  $D^{*0}\ell$  component



**FEI Tag:** Centered signal peaks

# (FEI) Tagging- Using Signal Probability

- **Signal Probability:** Multivariate classifier output to select high-purity  $B_{tag}$  candidates. (NOT A TRUE PROBABILITY\*)
- **Selection Tuning:** Tighter cut  $\rightarrow$  higher purity, lower efficiency



Tighter Cut: SigProb  $> 0.08$